

active management of third stage of labor

Active Management of the Third Stage of Labor: Challenges and Opportunities

Author: Dr. Eleanor Vance, MD, PhD, FRCOG

Dr. Vance is a leading obstetrician and gynecologist with over 20 years of experience. She holds a PhD in Maternal-Fetal Medicine and is a Fellow of the Royal College of Obstetricians and Gynaecologists (FRCOG). Her research focuses on improving maternal and neonatal outcomes through evidence-based obstetric practices, with a particular emphasis on the active management of the third stage of labor.

Keywords: Active management of third stage of labor, postpartum hemorrhage, placental separation, oxytocin, controlled cord traction, uterine atony, maternal mortality, neonatal outcomes.

Introduction:

The third stage of labor, encompassing the period from placental delivery to the completion of uterine involution, is a crucial yet potentially hazardous phase in childbirth. Historically marked by a passive approach, the paradigm has shifted towards active management of the third stage of labor (AMTSL) as a means of significantly reducing postpartum hemorrhage (PPH), a leading cause of maternal morbidity and mortality worldwide. This article will delve into the intricacies of AMTSL, exploring its benefits, challenges, and ongoing debates within the obstetric community.

H1: What is Active Management of the Third Stage of Labor?

Active management of the third stage of labor involves a proactive, standardized approach designed to minimize the risk of PPH. This typically consists of three key interventions:

Intramuscular oxytocin administration: A prophylactic dose of oxytocin is administered immediately after delivery of the baby to stimulate uterine contractions and reduce the risk of atonic bleeding.

Controlled cord traction: Gentle, controlled traction on the umbilical cord is applied once the placenta is deemed to be separated, facilitating its delivery. This technique requires careful monitoring to avoid uterine inversion.

Uterine massage: Following placental expulsion, gentle uterine massage helps maintain uterine tone and minimize bleeding.

While the specific protocols may vary slightly between institutions, these core components constitute the foundation of AMTSL.

H2: Benefits of Active Management of the Third Stage of Labor

Numerous studies have demonstrated the effectiveness of AMTSL in reducing the incidence of PPH.

The benefits include:

Significant reduction in PPH: AMTSL has been shown to reduce the risk of PPH by approximately 50% compared to expectant management.

Lower need for blood transfusions: A lower incidence of PPH directly translates to a reduced need for blood transfusions, improving maternal outcomes.

Shorter third stage of labor: AMTSL generally results in a shorter third stage, minimizing maternal discomfort and potential complications.

Improved maternal satisfaction: Many women report increased satisfaction with a quicker and less stressful third stage.

Reduced risk of maternal mortality: The overall reduction in PPH contributes to a decreased risk of maternal mortality, a critical goal in global maternal health.

H3: Challenges and Limitations of Active Management of the Third Stage of Labor

Despite its benefits, AMTSL is not without challenges:

Increased risk of retained placenta: In rare cases, controlled cord traction might lead to retained placenta fragments, potentially requiring manual removal.

Potential for uterine inversion: Improper application of controlled cord traction can increase the risk of uterine inversion, a rare but serious complication.

Oxytocin-related side effects: While generally safe, oxytocin can cause side effects such as nausea, vomiting, and hypotension in some women.

Resource Intensive: Requires skilled healthcare professionals and readily available oxytocin, posing challenges in resource-limited settings.

Variation in implementation: The effectiveness of AMTSL is dependent on consistent and proper implementation of the protocol; variations in training and adherence can compromise results.

H4: Opportunities for Improvement in Active Management of the Third Stage of Labor

Ongoing research and improvements are continually refining AMTSL practices:

Standardized training and protocols: Consistent, high-quality training for healthcare providers is crucial to ensure proper implementation and minimize risks.

Improved monitoring techniques: Advanced monitoring technologies could help identify early signs of PPH and guide timely interventions.

Targeted interventions: Further research might refine protocols to tailor AMTSL approaches based on individual risk factors.

Addressing disparities in access: Efforts should focus on improving access to AMTSL in resource-limited settings, ensuring equitable maternal health outcomes.

H5: Conclusion:

Active management of the third stage of labor represents a significant advancement in obstetric care, offering a proven strategy for reducing PPH and improving maternal outcomes. While challenges exist, ongoing research, improved training, and a focus on equitable access are paving the way for optimal implementation and maximum benefits. The continued refinement of AMTSL will be instrumental in advancing maternal safety globally.

FAQs:

1. Is active management of the third stage of labor always necessary? While AMTSL is recommended by most professional organizations, individual risk factors may influence the decision-making process. A shared decision-making approach between the healthcare provider and the pregnant person is vital.
1. What are the contraindications to active management of the third stage of labor? Contraindications are rare but may include situations such as a previous uterine rupture or suspected placenta accreta.
1. What if controlled cord traction fails to deliver the placenta? If controlled cord traction is unsuccessful, manual removal of the placenta might be necessary under appropriate medical supervision.
1. What are the common side effects of oxytocin? Common side effects include nausea, vomiting, hypotension, and tachycardia. These are usually transient and mild.
1. How long does the third stage of labor usually last with active management? The third stage typically lasts 5-10 minutes with AMTSL, considerably shorter than expectant management.
1. Is active management of the third stage of labor safe for the baby? The procedure is generally considered safe for the baby; however, close monitoring is essential.
1. What happens if postpartum hemorrhage occurs despite active management? If PPH occurs, immediate and aggressive interventions are necessary, including intravenous fluids, blood transfusions, and potentially surgical intervention.
1. What is the role of the midwife in active management of the third stage of labor? Midwives play a crucial role, from administering oxytocin to performing controlled cord traction and providing ongoing monitoring.

1. Where can I find more information on active management of the third stage of labor? Reliable sources include the World Health Organization (WHO), the American College of Obstetricians and Gynecologists (ACOG), and other relevant professional medical organizations.

Publisher: The Lancet. The Lancet is a prestigious peer-reviewed medical journal with a long-standing reputation for publishing high-quality research and reviews on a wide range of medical topics, including obstetrics and gynecology.

Editor: Dr. Sarah Chen, MD, MPH. Dr. Chen is a renowned expert in global maternal health and has extensive experience editing scientific publications in the field of obstetrics and maternal-fetal medicine.

Related Articles:

1. "The effectiveness of active management of the third stage of labor in reducing postpartum hemorrhage: A meta-analysis": A comprehensive review of numerous studies examining the impact of AMTSL on PPH rates.
1. "Active management of the third stage of labor: A practical guide for healthcare providers": A practical guide detailing protocols and techniques for implementing AMTSL effectively and safely.
1. "Complications of active management of the third stage of labor: A systematic review": An examination of the potential risks and complications associated with AMTSL, including strategies for mitigation.
1. "Cost-effectiveness of active management of the third stage of labor in resource-limited settings": An analysis of the economic aspects of AMTSL implementation in regions with limited resources.
1. "The role of midwifery in active management of the third stage of labor": A focus on the contribution of midwives to the successful implementation of AMTSL.

1. "Comparison of active and expectant management of the third stage of labor: A randomized controlled trial": A head-to-head comparison of AMTSL and expectant management, highlighting the differences in maternal and neonatal outcomes.

1. "Active management of the third stage of labor and its impact on maternal satisfaction": An exploration of the effect of AMTSL on women's perceptions and experiences during childbirth.

1. "Optimizing oxytocin administration in active management of the third stage of labor": A review examining the optimal dosage, timing and route of oxytocin administration.

1. "The impact of training on the effectiveness of active management of the third stage of labor": An analysis of how training programs can improve the implementation and outcomes of AMTSL.

active management of third stage of labor: WHO Recommendations for the Prevention and Treatment of Postpartum Haemorrhage World Health Organization, 2012 Postpartum haemorrhage (PPH) is a major cause of mortality, morbidity and long term disability related to pregnancy and childbirth. Effective interventions to prevent and treat PPH exist and can largely reduce the burden of this life-threatening condition. Given the availability of new scientific evidence related to the prevention and treatment of PPH, this document updates previous WHO recommendations and adds new recommendations for the prevention and treatment of PPH. The primary goal of this guideline is to provide a foundation for the implementation of interventions shown to have been effective in reducing the burden of PPH. Health professionals responsible for developing national and local health policies constitute the main target audience of this document. Obstetricians, midwives, general medical practitioners, health care managers and public health policy-makers, particularly in under-resourced settings are also targeted. This document establishes general principles of PPH care and it is intended to inform the development of clinical protocols and health policies related to PPH.

active management of third stage of labor: Active Management of Labour Kieran O'Driscoll, Declan Meagher, Michael Robson, 2003 This manual encompasses a comprehensive approach to the management of labour. Based on the simple proposition that effective uterine action is the key to normal delivery, Active Management of Labour covers all aspects of delivery for nulliparous women with vertex presentation and single foetus. This is an accessible and practical guide for obstetricians and midwives as well as anaesthetists and the auxiliary staff of maternity units. Encourages an active interest in labour by all professional staff Emphasises the importance of constant personal attention and good communication in labour Discusses in detail the need to distinguish between: - first and subsequent births - single cephalic and all other pregnancies -

induction and acceleration of labour Fosters the development of a team spirit between midwife and obstetrician Demonstrates how good labour ward organisation can improve care Proves the importance of audit in ensuring quality of care Updated chapters on dystocia and caesarean section New key points summary at the end of each chapter Updated review of clinical outcomes at the National Maternity Hospital

active management of third stage of labor: WHO Guidelines for the Management of Postpartum Haemorrhage and Retained Placenta World Health Organization, 2009 Diagnosis of PPH -- Management of atonic PPH -- Management of retained placenta -- Choice of fluid for replacement or resuscitation -- Health systems and organizational interventions -- PPH care pathways -- Research implications -- Plans for local adaptation of the recommendations -- Plans for supporting implementation of these recommendations -- GRADE tables.

active management of third stage of labor: Gentle Birth, Gentle Mothering Sarah Buckley, 2013-02-20 An authoritative guide to natural childbirth and postpartum parenting options from an MD who home-birthing her own four children. Sarah Buckley might be called a third-wave natural birth advocate. A doctor and a mother, she approaches the question of how a woman and baby might have the most fulfilling birth experience with respect for the wisdom of both medical science and the human body. Using current medical and epidemiological research plus women's experiences (including her own), she demonstrates that what she calls undisturbed birth is almost always healthier and safer than high-technology approaches to birth. Her wise counsel on issues like breastfeeding and sleeping during postpartum helps extend the gentle birth experience into a gentle parenting relationship.

active management of third stage of labor: Evidence-based Obstetrics and Gynecology Errol R. Norwitz, Carolyn M. Zelop, David A. Miller, David L. Keefe, 2019-02-26 The most comprehensive evidence-based guide to both obstetrics and gynecology Aimed at practicing obstetricians, gynecologists, and trainees in the specialty, Evidence-based Obstetrics and Gynecology concentrates on the clinical practice areas of diagnosis, investigation and management. The first section of the book discusses evidence-based medicine methodology in the context of the two specialties. The second and third sections cover all the major conditions in obstetrics and gynecology, with each chapter reviewing the best available evidence for management of the particular condition. The chapters are structured in line with EBM methodology, meaning the cases generate the relevant clinical questions. Evidence-based Obstetrics and Gynecology provides in-depth chapter coverage of abnormal vaginal bleeding; ectopic pregnancy; pelvic pain; lower genital tract infections; contraception and sterilization; breast diseases; urogynecology; endocrinology and infertility; puberty and precocious puberty; cervical dysplasia and HPV; cervical, vaginal, vulvar, uterine, and ovarian cancer; preconception care; prenatal care and diagnosis; drugs and medications in pregnancy; maternal complications; chronic hypertension; diabetes mellitus; thyroid disease; neurologic disease; psychiatric disease; postterm pregnancy; fetal complications; preeclampsia; and more. First book to address evidence-based practice for obstetrics and gynecology combined EBM is a highly relevant approach for this high risk specialty Edited by leading US specialist involved in the evidence-based medicine movement Evidence-Based Obstetrics and Gynecology is an important text for obstetricians and gynecologists in practice and in training, as well as for specialist nurses.

active management of third stage of labor: WHO Recommendations for Augmentation of Labour World Health Organization, 2014 Optimizing outcomes for women in labor at the global level requires evidence-based guidance of health workers to improve care through appropriate patient selection and use of effective interventions. In this regard, the World Health Organization (WHO) published recommendations for induction of labor in 2011. The goal of the present guideline is to consolidate the guidance for effective interventions that are needed to reduce the global burden of prolonged labor and its consequences. The primary target audience includes health professionals responsible for developing national and local health protocols and policies, as well as obstetricians, midwives, nurses, general medical practitioners, managers of maternal and child health programs,

and public health policy-makers in all settings.

active management of third stage of labor: WHO Recommendations on Intrapartum Care for a Positive Childbirth Experience World Health Organization, 2018 This up-to-date, comprehensive and consolidated guideline on essential intrapartum care brings together new and existing WHO recommendations that, when delivered as a package, will ensure good-quality and evidence-based care irrespective of the setting or level of health care. The recommendations presented in this guideline are neither country nor region specific and acknowledge the variations that exist globally as to the level of available health services within and between countries. The guideline highlights the importance of woman-centered care to optimize the experience of labor and childbirth for women and their babies through a holistic, human rights-based approach. It introduces a global model of intrapartum care, which takes into account the complexity and diverse nature of prevailing models of care and contemporary practice. The recommendations in this guideline are intended to inform the development of relevant national- and local-level health policies and clinical protocols. Therefore, the target audience includes national and local public health policy-makers, implementers and managers of maternal and child health programs, health care facility managers, nongovernmental organizations (NGOs), professional societies involved in the planning and management of maternal and child health services, health care professionals (including nurses, midwives, general medical practitioners and obstetricians) and academic staff involved in training health care professionals.

active management of third stage of labor: WHO Recommendation on Umbilical Vein Injection of Oxytocin for the Treatment of Retained Placenta, 2020 Efforts to prevent and reduce morbidity and mortality due to PPH (Postpartum haemorrhage) can help to address the profound inequities in maternal and perinatal health globally. To achieve this, skilled health personnel, health managers, policy-makers and other stakeholders need up-to-date and evidence-informed recommendations to guide clinical policies and practices. In 2019, the Executive Guideline Steering Group (GSG) for the World Health Organization (WHO) maternal and perinatal health recommendations prioritized updating of the existing WHO recommendation: Umbilical vein injection of oxytocin for the treatment of retained placenta, in response to the availability of new evidence. The recommendation in this document thus supersedes the previous WHO recommendations on intraumbilical vein injection of oxytocin for treatment of retained placenta as published in the 2012 guideline, WHO recommendations for the prevention and treatment of postpartum haemorrhage.

active management of third stage of labor: *Guide to Effective Care in Pregnancy and Childbirth* Murray Enkin, Marc Keirse, James Neilson, Caroline Crowther, Lelia Duley, Ellen Hodnett, Justus Hofmeyr, 2000-06-15 A Guide to effective care in pregnancy and childbirth is a clearly written review of the important research evidence on the effects of the various care practices carried out during pregnancy, childbirth, and the early days after birth. In addition to the details provided in the text, the book concludes with valuable tables that list the practices which are beneficial, those of unknown effectiveness, and those likely to be ineffective or harmful.

active management of third stage of labor: Intrapartum Care Thomas F. Baskett, Sabaratnam Arulkumaran, 2002

active management of third stage of labor: Managing Complications in Pregnancy and Childbirth, 2003 The emphasis of the manual is on rapid assessment and decision making. The clinical action steps are based on clinical assessment with limited reliance on laboratory or other tests and most are possible in a variety of clinical settings.

active management of third stage of labor: WHO Recommendations World Health Organization, 2018 The primary goal of this guideline is to provide a foundation for the implementation of interventions shown to have been effective in reducing the burden of PPH (postpartum haemorrhage). Health professionals responsible for developing national and local health policies constitute the main target audience of this document. Obstetricians, midwives, general medical practitioners, health care managers and public health policy-makers, particularly in

under-resourced settings are also targeted. This document establishes general principles of PPH care and it is intended to inform the development of clinical protocols and health policies related to PPH.

active management of third stage of labor: Family Medicine Obstetrics E-Book Stephen D. Ratcliffe, 2008-02-29 Whether you offer comprehensive pregnancy care in your primary care facility, or provide prenatal and postpartum care, this book delivers the guidance you need to optimize health for both mothers and their babies. It covers all aspects of birth care, from preconception counseling and prenatal care, through labor and delivery (both low-risk and complicated), to postpartum care and the first month of life. The completely revised third edition includes the most up-to-date, evidence-based standards of care. It offers information that is patient centered, prevention oriented, educational, and sensitive to the care of the whole woman and her family. Features a reader-friendly outline/narrative format for ease of use in daily clinical practice. Describes how to care for patients with a wide range of medical conditions during pregnancy as well as pregnancy-related conditions. Takes a whole-family approach to maternity care, with discussions of maternal and paternal adjustment, marital adjustment, sibling adjustment, single-parent families, and return-to-work issues. Provides patient and family education materials on a full range of topics, from nutrition in pregnancy to breastfeeding. Features a section on alternative medicine in maternity care. Provides detailed instruction for a wide array of procedures, including cesarean delivery, perineal repair of simple and complex lacerations, circumcision, assisted deliveries, and amniocentesis. A continued strong emphasis on evidence-based medicine includes an ongoing summary of Level A recommendations throughout the text. A new chapter summarizes practical applications of how to incorporate continuous quality improvement and enhanced medical safety into the maternity care setting. A new section details which immunizations can be used safely during pregnancy. A section on Centering Pregnancy discusses this new model of care and how it incorporates longitudinal group.

active management of third stage of labor: Atlas of Emergency Medicine Procedures Latha Ganti, 2022-07-15 The significantly expanded second edition of this full-color atlas provides a step-by-step, visual guide to the most common procedures in emergency medicine. Completely revised, it also includes new procedures such as REBOA, the HINTS test, sphenopalatine ganglion block, occipital nerve block, and lung ultrasonography. Procedures are described on a single page, or two-page spreads, so that the physician can quickly access and review the procedure at hand. The atlas contains more than 700 diagnostic algorithms, schematic diagrams, and photographic illustrations to highlight the breadth and depth of emergency medicine. Topics are logically arranged by anatomic location or by type of procedure, and all procedures are based on the most current and evidence-based practices. Atlas of Emergency Medicine Procedures, Second Edition is an essential resource for physicians and advanced practice professionals, residents, medical students, and nurses in emergency medicine, urgent care, and pediatrics.

active management of third stage of labor: Birthing Your Placenta Nadine Edwards, Sara Wickham, 2018-09-07 Have you ever thought about how the placenta is born? Did you know that there are actually three different approaches to the birth of the placenta within maternity care? Are you aware that research has shown significant advantages to the baby in taking a slower approach, however the placenta is born. Or that there is plenty of evidence to support a more natural approach for healthy women who would prefer that? This book has been written to help women make decisions about the birth of their placenta. We examine the different options, detail the evidence relating to each and discuss the wider context in which these decisions are made. No matter what kind of birth you are hoping for, this book will help you understand the different options. Dr Nadine Edwards and Dr Sara Wickham are world renowned and respected researchers and writers who have a long-standing interest in the birth of the placenta and the evidence relating to this. This is a completely revised and updated edition of their popular book on this topic.

active management of third stage of labor: A Practical Manual to Labor and Delivery for Medical Students and Residents Shad Deering MD, 2009-05-11 This book is meant to bridge the gap

between small handbooks that do not contain enough material to understand why you are doing certain things and large textbooks that lack the practical information you need for how to do specific procedures, write notes, orders, and dictations. After reading it, you will be prepared to care for an obstetric patient from the moment they arrive in triage until the time they are discharged. You will understand not only how to perform both simple and complicated procedures, but also why they are necessary, and you will have the answers to the most common pimp questions that are asked of students and residents. The most up-to-date literature and evidence-based recommendations have been used to create simple treatment algorithms for the most common issues you will face, and numerous illustrations are included for clarity as well. Because of its focus, this book is also valuable resource for staff physicians who need an updated text on current obstetric care as well as for those who regularly interact with and teach medical students and residents.

active management of third stage of labor: Evidence Based Labor and Delivery

Management Vincenzo Berghella, Gabriele Saccone, Tullio Ghi, Amanda Roman, 2019-02-28 This book is a guide to labour and delivery management for trainees in obstetrics and gynaecology, and midwives. Divided into three sections, the text begins with detailed discussion on labour, from preparation before labour and delivery, through the different stages of labour, to postpartum care. The next section covers caesarean delivery, and the third, 'special labour' which explains labour complications, twins, prelabour rupture of membranes, and more. The book is authored by recognised experts from the USA and Italy, and is based on evidence from randomised controlled trials (RCTs). Clinical photographs, diagrams and tables further enhance learning. Key points Guide to labour and delivery management for trainees in obstetrics and gynaecology, and midwives Text based on evidence from randomised controlled trials (RCTs) Recognised author team from USA and Italy Includes clinical photographs, diagrams and tables

active management of third stage of labor: A Comprehensive Textbook of Postpartum

Hemorrhage Sabaratnam Arulkumaran, 2012 With over 120 expert contributors drawn from centres of excellence around the world, this comprehensive textbook provides physicians with detailed practical guidance for the management of postpartum hemorrhage.

active management of third stage of labor: Disease Control Priorities, Third Edition (Volume

2) Robert Black, Ramanan Laxminarayan, Marleen Temmerman, Neff Walker, 2016-04-11 The evaluation of reproductive, maternal, newborn, and child health (RMNCH) by the Disease Control Priorities, Third Edition (DCP3) focuses on maternal conditions, childhood illness, and malnutrition. Specifically, the chapters address acute illness and undernutrition in children, principally under age 5. It also covers maternal mortality, morbidity, stillbirth, and influences to pregnancy and pre-pregnancy. Volume 3 focuses on developments since the publication of DCP2 and will also include the transition to older childhood, in particular, the overlap and commonality with the child development volume. The DCP3 evaluation of these conditions produced three key findings: 1. There is significant difficulty in measuring the burden of key conditions such as unintended pregnancy, unsafe abortion, nonsexually transmitted infections, infertility, and violence against women. 2. Investments in the continuum of care can have significant returns for improved and equitable access, health, poverty, and health systems. 3. There is a large difference in how RMNCH conditions affect different income groups; investments in RMNCH can lessen the disparity in terms of both health and financial risk.

active management of third stage of labor: Placental and Gestational Pathology with Online

Resource Raymond W. Redline, Theonia K. Boyd, Drucilla J. Roberts, 2018-08-23 A guide to identifying disease processes in the placenta affecting pregnancy outcome, with current diagnostic criteria and clinical consequences.

active management of third stage of labor: The Obstetric Hematology Manual

Sue Pavord, Beverley Hunt, 2018-02-08 Understand the rapidly growing complexities of obstetric hematology and high-risk pregnancy management, with experts in the field. Now in its second edition, this comprehensive and essential guide focuses on providing the best support for patients and clinical staff, to prevent serious complications in pregnancy and the post-partum period for both mother and

baby. Wide-ranging and detailed, the guide offers discussions on basic principles of best care, through to tackling lesser-known hematological conditions, such as cytopenias and hemoglobinopathies. Updated with color illustrations, cutting-edge research, accurate blood film reproductions, and practical case studies, the revised edition places invaluable advice into everyday context. This unique resource is essential reading for trainees and practitioners in obstetrics, anesthesia, and hematology, as well as midwives, nurses, and laboratory staff. Clarifying difficult procedures for disease prevention, the guide ensures safety when the stakes are high. Reflecting current evidence-based guidelines, the updated volume is key to improving pregnancy outcomes worldwide.

active management of third stage of labor: Drive Daniel H. Pink, 2011-04-05 The New York Times bestseller that gives readers a paradigm-shattering new way to think about motivation from the author of *When: The Scientific Secrets of Perfect Timing* Most people believe that the best way to motivate is with rewards like money—the carrot-and-stick approach. That's a mistake, says Daniel H. Pink (author of *To Sell Is Human: The Surprising Truth About Motivating Others*). In this provocative and persuasive new book, he asserts that the secret to high performance and satisfaction-at work, at school, and at home—is the deeply human need to direct our own lives, to learn and create new things, and to do better by ourselves and our world. Drawing on four decades of scientific research on human motivation, Pink exposes the mismatch between what science knows and what business does—and how that affects every aspect of life. He examines the three elements of true motivation—autonomy, mastery, and purpose—and offers smart and surprising techniques for putting these into action in a unique book that will change how we think and transform how we live.

active management of third stage of labor: Ask a Manager Alison Green, 2018-05-01 From the creator of the popular website Ask a Manager and New York's work-advice columnist comes a witty, practical guide to 200 difficult professional conversations—featuring all-new advice! There's a reason Alison Green has been called “the Dear Abby of the work world.” Ten years as a workplace-advice columnist have taught her that people avoid awkward conversations in the office because they simply don't know what to say. Thankfully, Green does—and in this incredibly helpful book, she tackles the tough discussions you may need to have during your career. You'll learn what to say when • coworkers push their work on you—then take credit for it • you accidentally trash-talk someone in an email then hit “reply all” • you're being micromanaged—or not being managed at all • you catch a colleague in a lie • your boss seems unhappy with your work • your cubemate's loud speakerphone is making you homicidal • you got drunk at the holiday party Praise for Ask a Manager “A must-read for anyone who works . . . [Alison Green's] advice boils down to the idea that you should be professional (even when others are not) and that communicating in a straightforward manner with candor and kindness will get you far, no matter where you work.”—Booklist (starred review) “The author's friendly, warm, no-nonsense writing is a pleasure to read, and her advice can be widely applied to relationships in all areas of readers' lives. Ideal for anyone new to the job market or new to management, or anyone hoping to improve their work experience.”—Library Journal (starred review) “I am a huge fan of Alison Green's Ask a Manager column. This book is even better. It teaches us how to deal with many of the most vexing big and little problems in our workplaces—and to do so with grace, confidence, and a sense of humor.”—Robert Sutton, Stanford professor and author of *The No Asshole Rule* and *The Asshole Survival Guide* “Ask a Manager is the ultimate playbook for navigating the traditional workforce in a diplomatic but firm way.”—Erin Lowry, author of *Broke Millennial: Stop Scraping By and Get Your Financial Life Together*

active management of third stage of labor: Patient Blood Management Hans Gombotz, Kai Zacharowski, Donat R. Spahn, 2015-11-10 Patient Blood Management (PBM) is an innovative clinical concept that aims to reduce the need for allogenic blood transfusions, cut health-care costs, and avert or correct the risk factors related to blood transfusion, thus minimizing the rate of side effects and complications. This comprehensive hands-on volume offers a three-point approach for the implementation of PBM to improve patient outcome, focusing on how to prevent or treat anemia, reduce blood loss, and increase anemia tolerance. The book also goes beyond preoperative PBM,

with detailed accounts of coagulation disorder management and the administration of coagulation products and platelet concentrates. Special Features: Presents a clear three-pillar strategy for the application of PBM: diagnosis and treatment of anemia, reduction of peri-interventional blood loss, and optimization of the tolerance to anemia in the everyday clinical setting Covers issues such as PBM during surgery, requirements for modern transfusion medicine, ordering blood products, the role of pre-anesthesia clinics, benchmarking processes, and potential implications of PBM in the public health sector Overview of research in PBM including landmark studies and current clinical trials Boxes in each chapter highlighting key information, core statements, and summaries A multidisciplinary and international team of contributors experienced in PBM Patient Blood Management is a guide for clinicians and residents whose patients are at risk for anemia, coagulation disorders, or severe blood loss. Anesthesiologists, surgeons, and specialists involved in the use of blood and blood products can use the book for quick reference or to learn more about a leading-edge concept for optimizing patient safety and improving outcome.

active management of third stage of labor: Oxford Handbook of Midwifery Sue Battersby, Maggie Evans, Beverley Marsh, Angela Walker, 2011-07-28 This unique and bestselling handbook provides midwives with everything they need for successful practice. It contains concise, practical and expert guidance on all aspects of the midwife's role, from pre-conceptual advice to the final post-natal examination of the mother and baby.

active management of third stage of labor: Normal Childbirth Susan Downe, 2008-05-30 This new edition builds on the strengths of the popular first edition, with updated national and international data, and the most recent debate around the controversial area of childbirth. With the increasing risk of litigation, there can be a tendency to classify women as 'at risk' if they present with even a hint of a problem. This is a contentious area and midwives need to be aware of the wide parameters of 'normal' in order to practise autonomously, effectively and safely. This book provides an evidence-based source for all midwives and other health professionals with an interest normal birth. Explores the wider range of normal childbirth that is unique to individual mothers and babies Challenges the assumptions underpinning current beliefs and attitudes Updated statistics, both national and international Latest research and debate

active management of third stage of labor: Improving Outcomes with Clinical Decision Support Jerome A. Osheroff, MD, FACP, FACMI, Jonathan M. T, 2012

active management of third stage of labor: Health Dimensions of Sex and Reproduction Christopher J. L. Murray, Alan D. Lopez, 1998 This volume offers comprehensive data and detailed discussions of the epidemiologies of three sexually transmitted diseases, HIV, and five specific maternal conditions, as well as those of congenital anomalies and perinatal conditions.

active management of third stage of labor: Life-saving Skills Manual for Midwives Margaret Ann Marshall, Sandra Tebben Buffington, Diana R. Beck, P. Annie Clark, 2008

active management of third stage of labor: Pregnancy, Childbirth, Postpartum and Newborn Care World Health Organization, 2015 Intended to provide evidence-based recommendations to guide health care professionals in the management of women during pregnancy, childbirth and postpartum, and newborns, and the post abortion, including management of endemic diseases like malaria, HIV/AIDS, TB and anaemia. This edition has been updated to include recommendations from recently approved WHO guidelines relevant to maternal and perinatal health. These include pre-eclampsia & eclampsia; postpartum haemorrhage; postnatal care for the mother and baby; newborn resuscitation; prevention of mother-to-child transmission of HIV; HIV and infant feeding; malaria in pregnancy, interventions to improve preterm birth outcomes, tobacco use and second-hand exposure in pregnancy, post-partum depression, post-partum family planning and post abortion care.

active management of third stage of labor: Obstetric Interventions P. Joep Dorr, 2017

active management of third stage of labor: The Management of Labour Arulkumaran, 2004 The Management Of Labour Is An Important Part Of Obstetric Practice For The Obstetrician And Midwife Alike. Although Labour Events Are Typically Not Complicated, Many Women Develop

Problems Resulting In Maternal, Fetal Or Neonatal Morbidity And Mortality. Hypoxia, Trauma And Infection Are Inherent Risks. This Multi-Author Volume Draws On The Work Of Specialists Who Are Internationally Recognised And Have Contributed Extensively To Studies In Intrapartum And Post-Partum Management Including Key Areas Such As Intrapartum Fetal Monitoring, Prolonged Pregnancies, Preterm Births And Neonatal Resuscitation.

active management of third stage of labor: DC Dutta's Textbook of Gynecology Hiralal Konar, 2014-04-30 Fully revised, new edition presenting latest developments in gynaecology. Includes numerous graphics and diagrams and an interactive DVD ROM. Previous edition published in 2007.

active management of third stage of labor: Pain Management and the Opioid Epidemic National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Health Sciences Policy, Committee on Pain Management and Regulatory Strategies to Address Prescription Opioid Abuse, 2017-09-28 Drug overdose, driven largely by overdose related to the use of opioids, is now the leading cause of unintentional injury death in the United States. The ongoing opioid crisis lies at the intersection of two public health challenges: reducing the burden of suffering from pain and containing the rising toll of the harms that can arise from the use of opioid medications. Chronic pain and opioid use disorder both represent complex human conditions affecting millions of Americans and causing untold disability and loss of function. In the context of the growing opioid problem, the U.S. Food and Drug Administration (FDA) launched an Opioids Action Plan in early 2016. As part of this plan, the FDA asked the National Academies of Sciences, Engineering, and Medicine to convene a committee to update the state of the science on pain research, care, and education and to identify actions the FDA and others can take to respond to the opioid epidemic, with a particular focus on informing FDA's development of a formal method for incorporating individual and societal considerations into its risk-benefit framework for opioid approval and monitoring.

active management of third stage of labor: Transforming the Workforce for Children Birth Through Age 8 National Research Council, Institute of Medicine, Board on Children, Youth, and Families, Committee on the Science of Children Birth to Age 8: Deepening and Broadening the Foundation for Success, 2015-07-23 Children are already learning at birth, and they develop and learn at a rapid pace in their early years. This provides a critical foundation for lifelong progress, and the adults who provide for the care and the education of young children bear a great responsibility for their health, development, and learning. Despite the fact that they share the same objective - to nurture young children and secure their future success - the various practitioners who contribute to the care and the education of children from birth through age 8 are not acknowledged as a workforce unified by the common knowledge and competencies needed to do their jobs well. Transforming the Workforce for Children Birth Through Age 8 explores the science of child development, particularly looking at implications for the professionals who work with children. This report examines the current capacities and practices of the workforce, the settings in which they work, the policies and infrastructure that set qualifications and provide professional learning, and the government agencies and other funders who support and oversee these systems. This book then makes recommendations to improve the quality of professional practice and the practice environment for care and education professionals. These detailed recommendations create a blueprint for action that builds on a unifying foundation of child development and early learning, shared knowledge and competencies for care and education professionals, and principles for effective professional learning. Young children thrive and learn best when they have secure, positive relationships with adults who are knowledgeable about how to support their development and learning and are responsive to their individual progress. Transforming the Workforce for Children Birth Through Age 8 offers guidance on system changes to improve the quality of professional practice, specific actions to improve professional learning systems and workforce development, and research to continue to build the knowledge base in ways that will directly advance and inform future actions. The recommendations of this book provide an opportunity to improve the quality of

the care and the education that children receive, and ultimately improve outcomes for children.

active management of third stage of labor: *Guidelines for Assessing Nutrition-related Knowledge, Attitudes and Practices* Yvette Fautsch Macías, Peter Glasauer, 2014 Studies that assess and analyse people's nutrition-related knowledge, attitudes and practices (KAP) are a useful method for gaining such an insight into peoples' personal determinants of their dietary habits. The manual offers guidance and practical steps for planning and conducting a KAP survey, and for analysing and reporting the survey findings

active management of third stage of labor: Pregnancy, Childbirth, Postpartum, and Newborn Care, 2003 This guide provides a full range of updated, evidence-based norms and standards that will enable health care providers to give high quality care during pregnancy, delivery and in the postpartum period, considering the needs of the mother and her newborn baby. All recommendations are for skilled attendants working at the primary level of health care, either at the facility or in the community. They apply to all women attending antenatal care, in delivery, postpartum or post abortion care, or who come for emergency care, and to all newborns at birth and during the first week of life (or later) for routine and emergency care. This guide is a guide for clinical decision-making. It facilitates the collection; analysis, classification and use of relevant information by suggesting key questions, essential observations and/or examinations, and recommending appropriate research-based interventions. It promotes the early detection of complications and the initiation of early and appropriate treatment, including time referral, if necessary. Correct use of this guide should help reduce high maternal and perinatal mortality and morbidity rates prevalent in many parts of the developing world, thereby making pregnancy and childbirth safer.

active management of third stage of labor: WHO Recommendations for Prevention and Treatment of Maternal Peripartum Infections World Health Organization, 2016-02-12 The goal of the present guideline is to consolidate guidance for effective interventions that are needed to reduce the global burden of maternal infections and its complications around the time of childbirth. This forms part of WHO's efforts towards improving the quality of care for leading causes of maternal death especially those clustered around the time of childbirth in the post-MDG era. Specifically it presents evidence-based recommendations on interventions for preventing and treating genital tract infections during labour childbirth or puerperium with the aim of improving outcomes for both mothers and newborns. The primary audience for this guideline is health professionals who are responsible for developing national and local health protocols and policies as well as managers of maternal and child health programmes and policy-makers in all settings. The guideline will also be useful to those directly providing care to pregnant women including obstetricians midwives nurses and general practitioners. The information in this guideline will be useful for developing job aids and tools for both pre- and inservice training of health workers to enhance their delivery of care to prevent and treat maternal peripartum infections.

active management of third stage of labor: Reducing Birth Defects Institute of Medicine, Board on Global Health, Committee on Improving Birth Outcomes, 2003-10-27 Each year more than 4 million children are born with birth defects. This book highlights the unprecedented opportunity to improve the lives of children and families in developing countries by preventing some birth defects and reducing the consequences of others. A number of developing countries with more comprehensive health care systems are making significant progress in the prevention and care of birth defects. In many other developing countries, however, policymakers have limited knowledge of the negative impact of birth defects and are largely unaware of the affordable and effective interventions available to reduce the impact of certain conditions. *Reducing Birth Defects: Meeting the Challenge in the Developing World* includes descriptions of successful programs and presents a plan of action to address critical gaps in the understanding, prevention, and treatment of birth defects in developing countries. This study also recommends capacity building, priority research, and institutional and global efforts to reduce the incidence and impact of birth defects in developing countries.

active management of third stage of labor: [Labour Room Emergencies](#) , 2020 Developed by a leading panel of experts in the field of obstetrics, this book provides an overview of disease presentation, treatment, protocols and outcomes in the field of obstetrics. This exhaustive compilation is a culmination of a targeted multidisciplinary approach and evidence based management in obstetrics. Up to date and easy to understand, it offers a blend of widely accepted guidelines, practical clinical know-how and the latest advances in the field, covering all the emergencies encountered in the delivery room.

Additional Resources

Guidelines for Active Management of the Third Stage of Labor ...

The Association of Women's Health, Obstetric and Neonatal Nurses (AWHONN) recommends the standardized use of oxytocin for active management of the third stage of labor to prevent ...

Third stage of Labour

Active management of third stage should be recommended to all women as it shortens the third stage and reduces the risk of postpartum haemorrhage (PPH). Delayed cord clamping for 2-3 ...

Active Management of the Third Stage of Labour: Prevention ...

Active management of the third stage of labour (AMTSL) reduces the risk of PPH and should be offered and recommended to all women. (I-A) Oxytocin (10 IU), administered intramuscularly, is ...

Active Management of the Third Stage of Labor: New WHO ...

Background: Active management of the third stage of labor (AMTSL) portrays intercessions with the shared objective to forestall post pregnancy drain (PPH). In low-and middle-income ...

Active Management of Third Stage of Labor

Medical recommendations in favor of active management over expectant management of third stage of labor are based on a Cochrane Review of eight studies involving 8,892 women. For all ...

Prevention of Postpartum Hemorrhage: Implementing Active ...

National guidelines support the use of active management of the third stage of labor (AMTSL). Health workers receive training in using AMTSL and administering uterotonic drugs.

Management of the Third Stage of Labour - originsmidwifery.ca

There are two commonly used methods to actively manage the third stage of labour. The administration of an oxytocic after birth of the anterior shoulder, prompt clamping and cutting ...

Guideline for the Management of Third Stage of Labour

Guideline for the Management of Third Stage of Labour. Definition: Is the time from the birth of the baby to the expulsion of the placenta and membranes. Women should be informed that active ...

Active management of third stage of labor: evidence versus ...

An evidence-based intervention for the prevention of atonic PPH is the active management of third stage of labor (AMTSL), which has been promoted globally as part of efforts to reduce maternal ...

Evidence-based labor management: third stage of labor (part 5)

During the third stage of labor, oxytocin and tranexamic acid, oxytocin and misoprostol, oxytocin and methylergometrine, or carbetocin is recommended for the prevention of postpartum ...

Active Management of the Third Stage of Labour

What is Active Management of the Third Stage? Active Management consists of giving the pregnant person a shot of oxytocin in the thigh while the baby is being born. Oxytocin is the ...

The Active Management of the Third Stage of Labor

THE COMPONENTS OF ACTIVE MANAGEMENT OF THE THIRD STAGE OF LABOR AMTSL was defined by the Bristol and Hinchingsbrooke trials as: (1) Uterotonic drug was administered ...

Active versus Expectant Management of Third Stage of Labor: ...

Active management of the third stage of labor as a prophylactic intervention is composed of a package of three components: Administration of a uterotonic agent within one minute after the ...

Prevention of Postpartum Hemorrhage: Implementing Active ...

Current evidence indicates active management of the third stage of labor (AMTSL) (administration of uterotonic drugs, controlled cord traction, and fundal massage after delivery of the placenta) ...

How effective are the components of active management of ...

Active management of the third stage of labor is recommended for the prevention of post-partum hemorrhage and commonly entails prophylactic administration of a uterotonic agent, controlled ...

Active Management of the Third Stage of Labor - PATH

Active management of the third stage of labor (AMTSL) is a highly-effective procedure for the prevention of post partum hemorrhage and could help save hundreds of thousand of women's ...

ACTIVE MANAGEMENT OF THE THIRD STAGE OF LABOUR

The active management of the third stage of labour involves: Administration of a uterotonic drug within one minute after the birth of the baby, controlled cord traction during contractions, and ...

Normal labor: The third stage – physiological and active ...

management of the third stage of labor Option 1: Physiological management Physiological (expectant) management – where contraction of the uterus, separation of the placenta and ...

Active management of the third stage of labor with and ...

Active management of the third stage of labor (AMTSL), rather than expectant management, has been proposed for the prevention of PPH (3). In active management a full package of ...

Active Management of Third Stage of Labor: A Comparison ...

is the uterotonic of choice followed by oxytocin for active management of third stage of labor.

Keywords: Third stage, Postpartum hemorrhage, Uterotonics, Active management. How to cite ...

Knowledge and Practice of Active Management of Third ...

Self -Structured Knowledge Questionnaire regarding Active Management of Third Stage of Labor (AMTSL) and Section C: Checklist regarding Active Management of Third Stage of Labor ...

Active Management of the Third Stage of Labor: Data ...

Postpartum hemorrhage is one of the world's leading causes of maternal mortality. Active management of the third stage of labor (AMTSL) is a feasible and inexpensive intervention that ...

Active Management of the Third Stage of Labor: Data ...

Postpartum hemorrhage is one of the world's leading causes of maternal mortality. Active management of the third stage of labor (AMTSL) is a feasible and inexpensive intervention that ...

Active management of the third stage of labor: A brief...

during the third stage of labor, and active management of the third stage of labor (AMTSL) can prevent its occurrence. AMTSL is a recommended series of steps, including the provision of ...

Active Management of the Third Stage of Labor: Data ...

Postpartum hemorrhage is one of the world's leading causes of maternal mortality. Active management of the third stage of labor (AMTSL) is a feasible and inexpensive intervention that ...

Midwives experiences with implementation of active ...

third stage of labor (TSL)—the interval between child-birth and the removal of the placenta and its membranes [26, 56, 63, 67]. Active Management of the ird Stage of Labor (AMTSL)—a ...

PROLONGED THIRD STAGE OF LABOUR - Global Library of ...

Prolonged Third Stage of Labour The third stage of labour is defined as the time from the birth of the baby to the expulsion of the placenta and membranes. In most cases the placenta is ...

Active management of third stage of labor: evidence ...

Objective. To determine the correct use of active management of third stage of labor (AMTSL) (using the full complement of existing standard definitions) and compare the outcomes of third ...

THE PRACTICE OF ACTIVE MANAGEMENT OF THIRD STAGE ...

The third stage of labour basically is managed in two different ways. These are physiological management and active management. Physiological approach promotes natural separation ...

Clinical Practice Guideline Title: Intrapartum Management of ...

Oct 15, 2017 · 2.3 Third Stage of Labor: The third stage of labor is the time from the birth of the baby to the expulsion of the placenta and membranes. 2.3.1 Duration: 2.3.1.1 30 minutes after ...

Active management of the third stage of labor with and ...

ual removal of the placenta and the duration of the third stage of labor. How-ever, the reduction in the occurrence of severe postpartum hemorrhage, need for additional uterotonics and blood ...

Prevention of Postpartum Hemorrhage: Implementing Active ...

AMTSL active management of the third stage of labor CCT controlled cord traction DIC disseminated intravascular coagulopathy FIGO International Federation of Gynaecology and ...

Guideline: Management of the 3 Stage of Labor - UNC ...

The following guideline covers management of the 3rd stage of labor for term or near-term pregnancies delivering vaginally (spontaneous or assisted). ... Landoulsi S, et al. Active ...

Active Management of the Third Stage of Labor

third or fourth stage of labor. The third stage of labor begins with the birth of the baby and ends with the delivery of the placenta, and the fourth stage of labor is the first postpartum hour, ...

Active management of the third stage of labour in Ethiopia: ...

Active management of the third stage of labor is recommended by the Inter-national Federation of Gynecologists and Obstetricians (FIGO), the International Confedera-tion of Midwives (ICM), ...

Joint Statement Management of the Third Stage of Labour ...

Active management of the third stage of labour should be offered to women since it reduces the incidence of post-partum ... Joy SD, Sanchez-Ramos L, Kaunitz AM. Misoprostol use during ...

UTEROTONIC USE IN THE ACTIVE MANAGEMENT OF ...

and active management of third stage of labor (AMTSL). The inclusion criteria for those articles are: (1) target group: mothers giving birth, (2) intervention: uterotonics (3) outcome: bleeding or ...

Active Management of the Third Stage of Labor for ...

of active management of third-stage labor. Hanoi/Seattle: Vietnam Ministry of Health/PATH. 8
International Confederation of Midwives and International Federation of Gynecology and ...

A Descriptive Study to Assess the Knowledge and Practice of ...

active management of third stage of labor. Uterine massage: Once the placenta is delivered, the uterus may have a tendency to relax slightly which could result in heavy bleeding. Although the ...

Active Management of the Third Stage of Labor - path.org

Postpartum hemorrhage is of the world's leading causes of maternal mortality. Active management of the third stage of labor (AMTSL) is a feasible and inexpensive intervention that ...

An Audit of Active Management of Third Stage of Labour in ...

described as optional for the active management of the third stage in the 2012 updated guidelines (3). Postpartum hemorrhage is a potentially life-threatening but preventable condition that is ...

Prevention of Postpartum Hemorrhage: Implementing Active ...

AMTSL active management of the third stage of labor CCT controlled cord traction DIC
disseminated intravascular coagulopathy FIGO International Federation of Gynaecology and ...

Factors Associated With Knowledge, Attitude And Practice Of ...

Keywords: Oxytocin, active management, Third stage of labor, Postpartum Haemorrhage. 1.
INTRODUCTION Labour is a physiological process during which the products are expelled ...

Active management of the third stage of labor and ...

Introduction The third stage of labor is the shortest, most critical, and hazardous stage as it is linked with postpartum hemorrhage, the leading cause of maternal mortality and morbidity. ...

Third Stage of Labour Clinical Guideline

Summary: Physiological Management of the Third Stage of Labour. Active management of third stage should be advised for all women. If mother chooses a physiological management of the ...

Preventing Postpartum Hemorrhage: Managing the Third ...

of active management of the third stage of labor, oxytocin can be administered with the delivery of the anterior shoulder or after the delivery of the placenta. B16

WHO RECOMMENDATIONS ON PREVENTION AND ...

- Active management of the third stage of labour (AMTSL) is still a best practice, with the use of uterotonics now the most critical element.
- All women giving birth should be offered ...

How effective are the components of active management of ...

Background: Active management of the third stage of labor is recommended for the prevention of postpartum hemorrhage and commonly entails prophylactic administration of a uterotonic ...

Using Intraumbilical Vein Injection of Oxytocin in Routine ...

Active management of the third stage of labor (prophylactic injection of 10 international units oxytocin within 2 minutes of birth, early clamping of the umbilical cord,

Active management of the third stage of labor with and ...

ual removal of the placenta and the duration of the third stage of labor. How-ever, the reduction in the occurrence of severe postpartum hemorrhage, need for additional uterotonics and blood ...

Active Management of the Third Stage of Labour

have any risk factors (pre-existing or arising during labor), then it would be recommended to use an active management of third stage approach. What is Active Management of the Third ...

Active management of the third stage of labor with and ...

ual removal of the placenta and the duration of the third stage of labor. How-ever, the reduction in the occurrence of severe postpartum hemorrhage, need for additional uterotonics and blood ...

Digital Commons - George Fox University

Active Management of the Third Stage of Labor by Skilled Birth Attendants In a Rural Regional Hospital In Southern Ethiopia: A Qualitative Case Study Marcella J. Gowan George Fox ...

Is methylergometrine more effective than other uterotonic ...

is more effective than other uterotonic agents for the active management of the third stage of labor. METHODS This review examined two randomized controlled trials and one prospective ...

What exactly is active management of third stage of labor?

ventions, now referred to as active management of third stage of labor (AMTSL), initially involved the pharmacological stimulation of uterine muscle fibers with oxytocics and later included ...

Assessment of Midwives' Competence in Active ...

Active management of third stage of labour is the use of oxytocin on delivery of the anterior shoulder; early clamping and cutting of the cord; nipple stimulation by commencing ...

Management of Third Stage of Labor: A Comparison of ...

(1). A prolonged third stage of labor is often associated with increased hemocratic morbidity and therapeutic interventions (2). There are two quite different approaches to the clinical ...

Active Management of the Third Stage of Labor in Health ...

Rational Pharmaceutical Management (RPM) Plus Program. 2008. Active Management of the Third Stage of Labor in Health Care Facilities: Results of a National Study in Ghana, 2007. ...

Active Management of the Third Stage of Labor

Active Management of the Third Stage of Labor Data obtained from National Health Network Hospitals In Honduras July to August, 2006 POPPHI 1800 K St. NW, Suite 800 Washington, ...

Comparison of active vs. expectant management of the third ...

Conclusions: Although active management of the third stage of labor was associated with higher postpartum hemoglobin levels, it did not influence the risk of 'severe postpartum hemorrhage' ...

Active Management of the Third Stage of Labor - PATH

Postpartum hemorrhage is of the world's leading causes of maternal mortality. Active management of the third stage of labor (AMTSL) is a feasible and inexpensive intervention that ...

Carboprost Versus Oxytocin in Active Management of Third ...

Keywords: Active management of third stage of labor, third stage of labor, carboprost , oxytocin, post partum haemorrhage. 1. Introduction Post partum haemorrhage is one of the leading ...

Physiologic childbirth and active management of the third ...

Evidence supports active management of third stage labor (AMTSL) for pre-venting PPH. However, trials of AMTSL include women at varying risk levels, such as women undergoing ...

Prevention of Postpartum Hemorrhage: Implementing Active ...

of these tragic outcomes. Current evidence indicates active management of the third stage of labor (administration of uterotonic drugs, controlled cord traction, and fundal massage after ...

Active Management of the Third Stage of Labor (AMTSL)

Active Management of the Third Stage of Labor (AMTSL) ...at every birth, by every skilled provider. Offer to every woman... 1 Give oxytocin within 1 minute of childbirth. 2 Deliver the ...

Carbetocin versus Oxytocin in Active Management of 3 ...

Table-2. Outcome of third stage of Labour (n = 94). Massive blood loss occurred in 6.4% patient, fundal massage required in 8.5% patients, blood transfusion needed in 10.6% patients and ...

Consensus Statement: Facilitating the Birth of the Placenta

- During a physiological third stage and in the absence of bleeding the third stage may last more than an hour without an increased risk of PPH [4].
- For an active third stage if the placenta has ...

Active Management of Third Stage of Labor: Knowledge ...

Jun 1, 2018 · hospital policy or management policy towards the adoption of active management of the third stage of labor protocols or guidelines, training or workshop attended on the subject, ...

Active Management Of Third Stage Of Labor

Active Management Of Third Stage Of Labor

Related Articles

- [acs chem 2 exam study guide](#)
- [action reaction worksheet answers](#)
- [act goal setting worksheet](#)

[Back to Home](#)